

CP Violation in Hadronic τ Decays

Alakabha Datta ^{a,1}, Ken Kiers ^{b,2}, David London ^{c,3},
Patrick J. O'Donnell ^{d,4}, and Alejandro Szynkman ^{c,5}

- a: Dept of Physics and Astronomy, 108 Lewis Hall,
University of Mississippi, Oxford, MS 38677-1848, USA*
*b: Physics Department, Taylor University,
236 West Reade Ave., Upland, Indiana, 46989, USA*
*c: Physique des Particules, Université de Montréal,
C.P. 6128, succ. centre-ville, Montréal, QC, Canada H3C 3J7*
*d: Department of Physics, University of Toronto,
60 St. George Street, Toronto, ON, Canada M5S 1A7*

(February 2, 2008)

Abstract

We re-examine CP violation in the $\Delta S = 0$ decays $\tau \rightarrow N\pi\nu_\tau$ ($N = 2, 3, 4$). We assume that the new physics (NP) is a charged Higgs. We show that there is no NP contribution to $\tau \rightarrow \pi\pi\nu_\tau$, which means that no CP violation is expected in this decay. On the other hand, NP can contribute to $\tau \rightarrow N\pi\nu_\tau$ ($N = 3, 4$). These are dominated by the intermediate resonant decays $\tau \rightarrow \omega\pi\nu_\tau$, $\tau \rightarrow \rho\pi\nu_\tau$ and $\tau \rightarrow a_1\pi\nu_\tau$. We show that the only sizeable CP-violating effects which are possible are in $\tau \rightarrow a_1\pi\nu_\tau \rightarrow 4\pi\nu_\tau$ (polarization-dependent rate asymmetry) and $\tau \rightarrow \omega\pi\nu_\tau$ (triple-product asymmetry).

¹datta@phy.olemiss.edu

²knkiers@taylor.edu

³london@lps.umontreal.ca

⁴odonnell@physics.utoronto.ca

⁵szynkman@lps.umontreal.ca

1 Introduction

One of the most important problems in particle physics is finding the origin of CP violation. CP violation was first seen in the kaon system about 40 years ago and was observed in B decays some 5 years ago [1]. If one hopes to understand CP violation, it is important to look for its effects in as many decays as possible. One of the purposes of the present paper is to push for the search for CP violation in hadronic τ decays. This is not a new area of research – there is already a large literature on the subject [2, 3, 4, 5, 6, 7, 8, 9, 10, 11]. However, there have been few experimental results lately, and many studies have focused on decays which are less likely to exhibit CP violation such as $\tau \rightarrow \pi\pi\nu_\tau$ [12].

All CP-violating effects come about due to the interference of (at least) two amplitudes with a relative weak (CP-odd) phase. In the standard model (SM) all τ decays involve (essentially) only a virtual W . To the extent that there is only one decay amplitude, all CP violation in τ decays vanishes in the SM. That is, the observation of CP violation in τ decays would be a smoking-gun signal of physics beyond the SM. (Note that there can be a SM CP asymmetry in $\tau \rightarrow \pi K_s \nu_\tau$ due to the several amplitudes in the K_s . The CP asymmetry is $O(10^{-3})$, see Ref. [13].)

Such CP violation will necessarily involve the interference of the SM W diagram with some new-physics (NP) amplitude. This NP amplitude typically consists of a right-handed W_R or a charged-Higgs exchange. We show that W - W_R interference always leads to effects which are negligible (they are suppressed by m_ν). Thus we need only consider a NP charged-Higgs exchange. This assumption, while not completely model-independent (e.g. we do not consider light leptoquarks), is fairly general – a great many models of NP include a light charged Higgs boson.

This said, many such models have two Higgs doublets, which give mass to the fermions. In such models, the coupling of the charged Higgs boson to first-generation final states is generally proportional to m_u and m_d , which are tiny. As a result, any CP violation which is due to charged-Higgs exchange is correspondingly small. If CP violation is to be observed in τ decays, the charged-Higgs coupling must be large. Thus, τ decays probe non-“standard” NP CP violation. This is arguably less likely to be observed, but if seen, the payoff is huge.

In this paper we focus on the decays $\tau \rightarrow N\pi\nu_\tau$ ($N = 2, 3, 4$). We first show that the charged Higgs cannot contribute to $\tau \rightarrow \pi\pi\nu_\tau$, so that no CP violation is expected in these decays. By contrast, the charged Higgs can contribute to decays with three and four pions, so that CP violation is possible. These decays are dominated by intermediate resonances (mostly vector). A second purpose of the present paper is to argue that, in studying $\tau \rightarrow N\pi\nu_\tau$ ($N = 3, 4$), it is best to concentrate on CP violation in the resonant decays. To this end, we present a study of CP-violating effects in $\tau \rightarrow V\pi\nu_\tau$ ($V = \omega, \rho, a_1$). We find that the total rate asymmetries in all decays are tiny. The polarization-dependent rate asymmetry may be measurable in $\tau \rightarrow a_1\pi\nu_\tau$. A triple-product asymmetry may be measured in $\tau \rightarrow \omega\pi\nu_\tau$. This

involves the measurement of the spin of the ω , which in turn implies knowledge of the τ momentum. We do not expect to measure large CP-violating effects in $\tau \rightarrow \rho\pi\nu_\tau \rightarrow 3\pi\nu_\tau$.

Sec. 2 contains a proof that W - W_R CP-violating effects are negligible, which leads us to consider charged-Higgs exchange as the new physics. We then discuss three possible CP-violating effects that are allowed, in general: the total rate asymmetry, the polarization-dependent rate asymmetry, and the triple-product asymmetry. We then examine the charged-Higgs contribution to $\tau \rightarrow N\pi\nu_\tau$ ($N = 2, 3, 4$). Our main results are found in Sec. 3. We consider resonant contributions to the hadronic τ decays and examine $\tau \rightarrow \omega\pi\nu_\tau$, $\tau \rightarrow \rho\pi\nu_\tau$ and $\tau \rightarrow a_1\pi\nu_\tau$. We write the form factors for the decays $\tau \rightarrow V\pi\nu_\tau$ ($V = \omega, \rho, a_1$) and examine the possible CP-violating effects. We find that such effects are possible in $\tau \rightarrow a_1\pi\nu_\tau \rightarrow 4\pi\nu_\tau$ (polarization-dependent rate asymmetry) and $\tau \rightarrow \omega\pi\nu_\tau$ (triple-product asymmetry). We conclude in Sec. 4. Several technical details are included in the Appendix.

2 τ Decays: Generalities

2.1 SM-NP Interference

Consider the decay $\tau \rightarrow f\nu_\tau$, where f is some final hadronic state. Assume also that the SM and NP contribute to this decay. Both contributions can be split into a leptonic and a hadronic piece. The SM leptonic piece is proportional to $\bar{u}_\nu\gamma_\mu\gamma_L u_\tau$, where $\gamma_L = (1 - \gamma_5)/2$.

Now suppose that the NP consists of a right-handed W_R . In this case, the NP leptonic piece is proportional to $\bar{u}_\nu\gamma_\mu\gamma_R u_\tau$, where $\gamma_R = (1 + \gamma_5)/2$. The SM-NP interference thus includes a piece proportional to $\text{Tr}[u_\nu\bar{u}_\nu\gamma_\mu\gamma_L u_\tau\bar{u}_\tau\gamma_\nu\gamma_R]$. This trace is proportional to m_ν : $\gamma_R u_\nu\bar{u}_\nu\gamma_R \sim m_\nu$. Since m_ν is negligible, we therefore conclude that CP violation in τ decays coming from W - W_R interference is negligible, and take the NP to consist only of a charged Higgs boson H .

The coupling of the charged Higgs to leptons is scalar and/or pseudoscalar. This coupling can be written in terms of γ_L and γ_R . The piece of the neutrino coupling proportional to γ_R leads to a W - H interference term which is proportional to m_ν (as in the interference of W and W_R above), and is negligible. Thus, it is only the H -neutrino coupling proportional to γ_L which is important. The W - H interference term then involves

$$\begin{aligned} \text{Tr}[u_\nu\bar{u}_\nu\gamma_\mu\gamma_L u_\tau\bar{u}_\tau\gamma_\nu] &= \text{Tr}\left[\not{p}_\nu\gamma_\mu\gamma_L(\not{p}_\tau + m_\tau)\frac{(1 + \gamma_5\not{s}_\tau)}{2}\gamma_L\right] \\ &= \frac{1}{2}m_\tau\text{Tr}[\not{p}_\nu\gamma_\mu\gamma_L] + \frac{1}{2}\text{Tr}[\not{p}_\nu\gamma_\mu\not{p}_\tau\not{s}_\tau\gamma_L]. \end{aligned} \quad (1)$$

The first term is always present; the second term is only important if the spin of the τ is known (i.e. if the τ is polarized).

2.2 CP-Violating Effects

If there are two contributing amplitudes to the decay $\tau \rightarrow f\nu_\tau$ then the full decay amplitude can be written

$$\mathcal{A} = A_1 + A_2 e^{i\phi} e^{i\delta}, \quad (2)$$

where ϕ and δ are the relative weak (CP-odd) and strong (CP-even) phases, respectively. The full rate is proportional to $\sum_{spins} |\mathcal{A}|^2$. If some spins are measured, one does not sum over them.

One CP-violating signal occurs if the spin-independent or spin-dependent rate is different from that of its anti-process. This is known as direct CP violation. It is proportional to

$$\sin \phi \sin \delta. \quad (3)$$

Thus, one can obtain a direct CP asymmetry if the two decay amplitudes have a nonzero relative weak *and* strong phase.

A second signal involves a *triple product* (TP). The TP takes the form $\vec{v}_1 \cdot (\vec{v}_2 \times \vec{v}_3)$, where each v_i is a spin or momentum. In a 2-body τ decay, the TP must involve 2 spins – since the ν_τ cannot be measured, this must include the spin of the τ . The TP for a 3-body decay includes 1 spin; all the v_i can be momenta in an M -body ($M \geq 4$) decay. If the TP in a given τ decay is different from that seen in its anti-process, this is another signal of CP violation. The TP asymmetry is proportional to

$$\sin \phi \cos \delta. \quad (4)$$

Thus, one does not require a strong-phase difference to get a TP asymmetry.

As discussed above, all CP-violating effects require the interference of two amplitudes. In particular, the transition $H \rightarrow f$ must be possible. For example, suppose that f can be produced only through the decay of a vector (or axial-vector) meson V . Since the spin-0 charged Higgs cannot couple to the spin-1 vector meson, the decay $\tau \rightarrow V\nu_\tau \rightarrow f\nu_\tau$ can come only from a SM virtual W exchange. In this case there is only a single amplitude and one will not have CP violation. The lesson here is that one must check whether the final-state f can be produced from a spin-0 charged Higgs boson.

2.3 $\Delta S = 0$ τ Decays: $\tau \rightarrow N\pi\nu_\tau$

Above, we argued that, in the decay $\tau \rightarrow f\nu_\tau$, it is crucial to know if the transition $H \rightarrow f$ is allowed. Here we consider $\tau \rightarrow N\pi\nu_\tau$, with $N = 2, 3, 4$. For a given $N\pi$ final hadronic state one can use isospin to determine if $H \rightarrow f$ can take place.

Consider $N = 2$, i.e. $\tau \rightarrow \pi\pi\nu_\tau$. The intermediate particle (W, H) decays to $\bar{u}d$. The $\pi\pi$ final state can be produced if a gluon decays to a $u\bar{u}$ or $d\bar{d}$ pair, and these quarks combine with the $\bar{u}d$ from the decaying particle. In the isospin limit, the $u\bar{u}$ and $d\bar{d}$ quark pairs have the same coupling to the gluons, and are produced with the same rate. The amplitude for $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ thus involves the formation of mesons

from quarks. Under isospin we have $\pi^+ \equiv \bar{d}u$, $\pi^- \equiv -\bar{u}d$ and $\pi^0 \equiv (\bar{d}d - \bar{u}u)/\sqrt{2}$. Thus,

$$\bar{u}(u\bar{u})d + \bar{u}(d\bar{d})d \sim [\pi^0\pi^- - \pi^-\pi^0] . \quad (5)$$

A priori, the state of two pions corresponds to total isospin $I = 0, 1, 2$. However, this antisymmetric combination corresponds to $I = 1$ only. (Note: the total isospin must be $I = 1$ because the gluons, which produce the quark pairs, have $I = 0$.) Since the total wave function of two pions must be symmetric (Bose symmetry), the relative orbital angular momentum must be odd, giving a spatial wave function which is also antisymmetric. This says that the $\pi\pi$ final state *cannot* be produced from a spin-0 charged Higgs boson. Thus, no CP violation is expected in this decay (more precisely, it is only at the level of isospin breaking). This is what is found experimentally [12].

Note that the above argument does not hold when the final state A^-B^0 does not involve identical particles. Examples of this are $\rho^0\pi^-$, $a_1^-\pi^0$, $\omega^0\pi^-$, etc. That is, we know that the final state must have $I = 1$, but this does not imply anything about the total angular momentum in the absence of identical particles. Thus, $J = 0$ is allowed, and the final state A^-B^0 can be produced in the decay of a charged Higgs. For these final states, CP-violating effects are expected.

A special case of this is the decay $\tau \rightarrow \pi K \nu_\tau$ which has been considered in Refs. [2, 3, 13, 14]. Here isospin gives $K^- \equiv -\bar{u}s$ and $\bar{K}^0 \equiv \bar{d}s$. Thus,

$$\bar{u}(u\bar{u})s + \bar{u}(d\bar{d})s \sim \left[(1/2)\pi^0 K^- - \pi^- \bar{K}^0 \right] . \quad (6)$$

Here the two pieces do not contain the same particles, and so the argument used above for $\tau \rightarrow \pi\pi\nu_\tau$ does not apply. Thus, the decay $\tau \rightarrow \pi K \nu_\tau$ receives (non-resonant) contributions from a charged Higgs boson. In addition, $\tau \rightarrow \pi K \nu_\tau$ can also be produced resonantly, from an intermediate vector V [$K^*(892)$] or an intermediate scalar S [$K_0^*(1430)$] [2]. Note that the charged Higgs can couple to an S . Thus CP violation is possible in $\tau \rightarrow \pi K \nu_\tau$. Since we have assumed that the coupling of the charged Higgs is arbitrary – that is, it is not proportional to quark masses – this CP violation could be large. However, it is also possible that the NP follows the pattern of the CKM matrix, so that CP violation in $\Delta S = 1$ decays is suppressed compared to that in $\Delta S = 0$ decays. In this paper we focus on $\Delta S = 0$ decays of the τ .

Now consider $N = 3$: $\tau \rightarrow \pi\pi\pi\nu_\tau$ [4, 5]. This actually represents two decays: $\tau^- \rightarrow \pi^-\pi^+\pi^-\nu_\tau$ (called $0\pi^0$) and $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ ($2\pi^0$). We write,

$$\begin{aligned} \pi\pi\pi &\sim \bar{u}(u\bar{u})(u\bar{u})d + \bar{u}(d\bar{d})(d\bar{d})d + \bar{u}(u\bar{u})(d\bar{d})d + \bar{u}(d\bar{d})(u\bar{u})d \\ &\sim -\pi^0\pi^0\pi^- - \pi^-\pi^0\pi^0 + \pi^0\pi^-\pi^0 + \pi^-\pi^+\pi^- . \end{aligned} \quad (7)$$

In this case there is no constraint on the total angular momentum of the $0\pi^0$ or $2\pi^0$ state, and the spin-0 H can contribute directly to both 3π τ decays in addition to the spin-1 W .

The $N = 4$ decay is similar: $\tau \rightarrow \pi\pi\pi\pi\nu_\tau$. As above, this actually represents two decays: $\tau^- \rightarrow \pi^-\pi^+\pi^-\pi^0\nu_\tau$ ($1\pi^0$) and $\tau^- \rightarrow \pi^-\pi^0\pi^0\pi^0\nu_\tau$ ($3\pi^0$). Given that the additional quarks come from gluons,

$$\begin{aligned}
\pi\pi\pi\pi &\sim \bar{u}(u\bar{u})(u\bar{u})(u\bar{u})d + \bar{u}(u\bar{u})(u\bar{u})(d\bar{d})d + \bar{u}(u\bar{u})(d\bar{d})(d\bar{d})d \\
&\quad + \bar{u}(d\bar{d})(d\bar{d})(d\bar{d})d + \bar{u}(d\bar{d})(u\bar{u})(d\bar{d})d + \bar{u}(d\bar{d})(u\bar{u})(u\bar{u})d \\
&\quad + \bar{u}(d\bar{d})(d\bar{d})(u\bar{u})d + \bar{u}(u\bar{u})(d\bar{d})(u\bar{u})d \\
&\sim \pi^0\pi^0\pi^0\pi^- - \pi^0\pi^0\pi^-\pi^0 + \pi^0\pi^-\pi^0\pi^0 - \pi^-\pi^0\pi^0\pi^0 \\
&\quad + \pi^-\pi^+\pi^-\pi^0 - \pi^-\pi^+\pi^0\pi^- + \pi^-\pi^0\pi^+\pi^- - \pi^0\pi^-\pi^+\pi^- . \quad (8)
\end{aligned}$$

The first and last four terms contribute to the decays $3\pi^0$ and $1\pi^0$, respectively. These terms do not have any particular symmetry under the exchange of any two pairs of π 's, so that there is no constraint on the total angular momentum of the final state. Thus, the spin-0 H can contribute directly (non-resonantly) to both 4π τ decays.

3 $\tau \rightarrow \omega\pi\nu_\tau / \rho\pi\nu_\tau / a_1\pi\nu_\tau$

We have therefore seen that there can be non-resonant contributions to $\tau \rightarrow N\pi\nu_\tau$ ($N = 3, 4$) coming from the charged Higgs. However, there can also be resonant contributions from intermediate states such as $\omega\pi$, $\rho\pi$ and $a_1\pi$. These are 3-body decays, while the non-resonant decays are 4-body or 5-body. The phase-space suppression of these latter decays implies that the widths of such decays are much smaller than those of resonant decays, and so $\tau \rightarrow N\pi\nu_\tau$ is dominated by resonances. (This is consistent with the results of Ref. [15], which finds that the 4π 's in $\tau \rightarrow \pi\pi\pi\pi\nu_\tau$ come primarily from the (resonant) decay of $\omega\pi$ or $a_1\pi$.) In the following we neglect all non-resonant contributions.

Thus, there are several $\Delta S = 0$ τ decays which can receive contributions from a spin-0 charged Higgs. That is, charged H (NP) can give rise to CP violation. Although the decay $\tau \rightarrow \pi\pi\nu_\tau$ will not manifest CP violation, such an effect can be seen in $\tau \rightarrow \omega\pi\nu_\tau$, $\tau \rightarrow \rho\pi\nu_\tau$, and $\tau \rightarrow a_1\pi\nu_\tau$. Obviously, the amount of CP violation will depend on the size of the $H \rightarrow f$ contribution.

In this section we examine CP violation in resonant $\Delta S = 0$ decays. There are two ingredients: the size of form factors (for both the SM and the NP), and the size of the hadronic charged Higgs coupling. These will be discussed in turn.

The matrix element for $\tau \rightarrow f\nu_\tau$ involves the product of a leptonic and hadronic current for the W and the charged Higgs. However, as we saw in Sec. 2.1, the Higgs' leptonic current must contain a left-handed coupling to neutrinos in order to interfere with that of the SM. We can therefore write the Hamiltonian as

$$H = \frac{G_F}{\sqrt{2}} \cos\theta_c (L_\mu H^\mu + L_0 H_0) + h.c. , \quad (9)$$

where the first and second pieces correspond to W and H exchange, respectively. We have

$$\begin{aligned} L_\mu &= \bar{\nu}\gamma_\mu(1 - \gamma_5)\tau \quad , \quad H^\mu = \bar{d}\gamma^\mu(1 - \gamma_5)u \\ L_0 &= \bar{\nu}(1 + \gamma_5)\tau \quad , \quad H_0 = \bar{d}(a + b\gamma_5)u . \end{aligned} \quad (10)$$

To evaluate the matrix element of $\tau \rightarrow f\nu_\tau$ we therefore have to estimate the hadronic matrix elements $\langle f|H^\mu|0\rangle$ and $\langle f|H_0|0\rangle$.

3.1 $\tau \rightarrow \omega\pi\nu_\tau/\rho\pi\nu_\tau/a_1\pi\nu_\tau$: Form Factors

Consider the decay $\tau(l) \rightarrow V(q_1)\pi(q_2)\nu_\tau(l')$, where V is a vector or axial-vector meson. The general structure for the SM current $J^\mu = \langle V(q_1)\pi(q_2)|H^\mu|0\rangle$ is [6, 7]

$$\begin{aligned} J^\mu &= F_1(Q^2) \left(Q^2 \epsilon_1^\mu - \epsilon_1 \cdot q_2 Q^\mu \right) + F_2(Q^2) \epsilon_1 \cdot q_2 \left(q_1^\mu - q_2^\mu - Q^\mu \frac{Q \cdot (q_1 - q_2)}{Q^2} \right) \\ &\quad + iF_3(Q^2) \varepsilon^{\mu\alpha\beta\gamma} \epsilon_{1\alpha} q_{1\beta} q_{2\gamma} + F_4(Q^2) \epsilon_1 \cdot q_2 Q^\mu , \end{aligned} \quad (11)$$

where $Q^\mu \equiv (q_1 + q_2)^\mu$ and ϵ_1 denotes the polarization tensor of the V . The latter satisfies the relations

$$q_1 \cdot \epsilon_1 = 0 , \quad (12)$$

$$\sum_\lambda \epsilon_1^\mu \epsilon_1^{*\nu} = -g^{\mu\nu} + \frac{q_1^\mu q_1^\nu}{m_V^2} . \quad (13)$$

The SM hadronic current is described by the four form factors $F_{1-4}(Q^2)$. For a vector meson (ω or ρ), experimentally the contribution proportional to F_3 dominates. At present there is no evidence for the “second-class currents” proportional to F_1 and F_2 [15, 16, 17] (which violate G -parity). For an axial-vector meson (a_1) F_3 switches roles with F_1 and F_2 . That is, for the a_1 , the contributions proportional to F_1 and F_2 dominate, while the F_3 term violates G -parity. In both cases the term proportional to F_4 can arise only from a scalar interaction. In the SM, this is negligible [7]. A significant F_4 term can only be produced from NP.

The current J^μ can receive pole contributions. For example, for $\omega\pi$ these come from the ρ and $\rho' = \rho(1450)$, which contribute to F_3 ,

$$F_3^{\text{pole}}(Q^2) = \frac{g_{\rho\omega\pi}}{\gamma_\rho} \left[\frac{m_\rho^2}{Q^2 - m_\rho^2 + i\sqrt{Q^2}\Gamma_\rho} + A_1 \frac{m_{\rho'}^2}{Q^2 - m_{\rho'}^2 + i\sqrt{Q^2}\Gamma_{\rho'}} \right] . \quad (14)$$

In this expression, $g_{\rho\omega\pi}$ is the $\rho\omega\pi$ strong coupling and γ_ρ is a quantity associated with the weak coupling of the ρ to the weak charged current [15]. The constant A_1 may be expressed as a ratio comparing $g_{\rho\omega\pi}$ and γ_ρ to the analogous quantities for the ρ' . In Ref. [15], A_1 is taken as a parameter that is fit to the data. It is

now clear that the widths of the ρ and the ρ' can generate a strong phase for the form factors. It should be pointed out that our model of the resonant contribution is simple and should be understood as a demonstration of how strong phases can arise. Note that, if G -parity is violated, there can also be resonant contributions to $F_1(Q^2)$ and $F_2(Q^2)$ [7].

The non-resonant part of J^μ may be estimated in Chiral Perturbation Theory [8] as an expansion in the pion momentum with the vector mesons treated as heavy. However, in τ decays the pion is not always soft and so higher-order terms in the chiral Lagrangian are important, leading to a loss of predictive power. Hence we do not estimate the non-resonant part.

We now turn to the charged Higgs' hadronic current (H_0). Since the charged Higgs has spin 0, the $V\pi$ system is in a relative $L = 1$ angular momentum state and so only the pseudoscalar (scalar) term in Eq. (10) can contribute to the hadronic current for a vector (axial-vector) meson V . The general structure is

$$J_{\text{Higgs}} = \langle V(q_1)\pi(q_2)|H_0|0\rangle = \begin{cases} bf_H \epsilon_1 \cdot q_2, & V: \text{ vector}, \\ af_H \epsilon_1 \cdot q_2, & V: \text{ axial-vector}, \end{cases} \quad (15)$$

where a and b are, respectively, the scalar and pseudoscalar couplings of the charged Higgs [Eq. (10)]. The form factor f_H can have a CP-conserving strong phase from rescattering while a and b will in general have a CP-violating weak phase from New Physics.

The squared matrix element is now given as

$$|\mathcal{A}|^2 = \frac{G_F^2}{2} \cos^2 \theta_c (M_\mu J^\mu + M_{\text{Higgs}} J_{\text{Higgs}}) (M_\nu J^\nu + M_{\text{Higgs}} J_{\text{Higgs}})^\dagger, \quad (16)$$

where the M 's are the matrix elements of the leptonic currents in Eq. (10). Using the equations of motion for the τ and the neutrino, we can rewrite Eq. (16) as

$$|\mathcal{A}|^2 = \frac{G_F^2}{2} \cos^2 \theta_c L_{\mu\nu} \widetilde{H}^{\mu\nu}, \quad (17)$$

where $L_{\mu\nu} \equiv M_\mu (M_\nu)^\dagger$ and $\widetilde{H}^{\mu\nu} \equiv \widetilde{J}^\mu (\widetilde{J}^\nu)^\dagger$, with

$$\widetilde{J}^\mu = \begin{cases} J^\mu + b(f_H/m_\tau)\epsilon_1 \cdot q_2 Q^\mu, & V: \text{ vector}, \\ J^\mu + a(f_H/m_\tau)\epsilon_1 \cdot q_2 Q^\mu, & V: \text{ axial-vector}. \end{cases} \quad (18)$$

Comparing Eqs. (11) and (18), we see that $\widetilde{J}^\mu$ is obtained from J^μ by the replacement $F_4(Q^2) \rightarrow \widetilde{F}_4(Q^2)$, where

$$\widetilde{F}_4(Q^2) = \begin{cases} F_4(Q^2) + (bf_H/m_\tau), & V: \text{ vector}, \\ F_4(Q^2) + (af_H/m_\tau), & V: \text{ axial-vector}. \end{cases} \quad (19)$$

Since any CP-violating effect requires an interference with the NP, we immediately see that $\widetilde{F}_4(Q^2)$ must be involved in any such effect.

For future reference we note that, in the hadronic rest frame (where $\vec{Q} = 0$ and $Q^0 = \sqrt{Q^2}$), we have

$$\tilde{J}^0 = \tilde{F}_4(Q^2) \epsilon_1 \cdot q_2 Q^0, \quad (20)$$

$$\tilde{J}^i = J^i = F_1(Q^2) Q^2 \epsilon_1^i + 2F_2(Q^2) \epsilon_1 \cdot q_2 q_1^i - iF_3(Q^2) \epsilon^{ij0k} \epsilon_{1j} q_{1k} Q_0. \quad (21)$$

Thus, in the hadronic rest frame, only $\tilde{J}^0$ contains a NP contribution.

Now for the decay $\tau^-(l) \rightarrow V(q_1)\pi^-(q_2)\nu_\tau(l')$ we can write the differential decay width as [1]

$$d\Gamma = \frac{(2\pi)^4}{2m_\tau} |\mathcal{A}|^2 d\Phi_3$$

$$d\Phi_3 = \delta^4(l - q_1 - q_2 - l') \frac{d^3q_1}{(2\pi)^3 2E_{q_1}} \frac{d^3q_2}{(2\pi)^3 2E_{q_2}} \frac{d^3l'}{(2\pi)^3 2E_{l'}}. \quad (22)$$

A similar differential distribution can be written for the CP conjugate process $\tau^+(l) \rightarrow V(q_1)\pi^+(q_2)\nu_\tau(l')$.

The differential distribution is best analyzed in the rest frame of the hadron ($\vec{q}_1 + \vec{q}_2 = 0$), as is done in Refs. [7, 9, 10]. (In the following we follow closely the conventions and notation of these references.) We define the angles α , β , θ and ψ in the same manner as they are defined in Ref. [7]. The definitions are reviewed here for convenience. Note that α , β and ψ are defined in the hadronic rest frame, while θ is defined in the τ rest frame.

Let the unit vector $\hat{n}_L$ denote the direction of the lab as viewed in the hadronic rest frame and let $\hat{q}_1 \equiv \vec{q}_1/|\vec{q}_1|$, where $\vec{q}_1$ is the V 's momentum in the hadronic rest frame. Then we let β denote the angle between $\hat{n}_L$ and $\hat{q}_1$, as indicated in Fig. 1. The angles α and ψ are also defined in Fig. 1. As noted above, θ is defined in the rest frame of the τ . In that frame, θ is the angle between the direction of flight of the τ in the laboratory frame and the direction of the hadrons. Note that in some cases, even if the τ 's direction cannot be determined experimentally, the angles θ and ψ can both be measured since they are related in a simple way to the energy E_h of the hadronic system [7].

We assume here that the τ 's have polarization P , aligned with their direction of flight in the laboratory. Then $s_\mu s^\mu = -P^2$. A derivation of the τ spin four-vector s^μ in the hadronic rest frame may be found in Appendix A of Ref. [9]. For convenience, an expression is included in the Appendix of the present work as well.

Since τ decays always involve a neutrino, it can be difficult in practice to reconstruct the τ 's momentum. At this stage, we assume that the τ direction is not known. Later, in our discussion of triple-product CP asymmetries, we will assume that the τ direction is in fact known. (Note that while the missing neutrino certainly makes the determination of the τ 's direction difficult, there are experimental techniques that can be employed to obtain the direction [11].)




Figure 1: Coordinate system showing the angles for the decay $\tau \rightarrow V\pi\nu$. The momentum vectors shown are in the hadronic rest frame. Note: this figure is very similar to one shown in Ref. [3].

Employing the various definitions for the angles, and integrating over α (the unknown τ direction), we find [7]

$$d\Gamma = \frac{|\vec{q}_1|}{2m_\tau(4\pi)^3} \left(\frac{m_\tau^2 - Q^2}{m_\tau^2} \right) \frac{dQ^2}{\sqrt{Q^2}} \frac{d\cos\theta}{2} \frac{d\cos\beta}{2} |\mathcal{A}|^2. \quad (23)$$

The matrix element squared, $|\mathcal{A}|^2$, is proportional to $L_{\mu\nu} \widetilde{H}^{\mu\nu}$ [see Eq. (17)]. We write the physical polarization states of the V in the hadronic rest frame as

$$\epsilon_1^\mu(\pm) = \frac{1}{\sqrt{2}} (0; \mp 1, i, 0), \quad \epsilon_1^\mu(0) = \left(-\frac{|\vec{q}_1^z|}{m_V}; 0, 0, -\frac{E_1}{m_V} \right), \quad (24)$$

where the coordinate system is defined in Fig. 1. In this basis, the hadronic tensor $\widetilde{H}^{\mu\nu} = \widetilde{J}^\mu (\widetilde{J}^\nu)^\dagger$ is simplified considerably [see Eqs. (20) and (21), and we sum over

polarizations:

$$\begin{aligned} \widetilde{H}^{\mu\nu} &\sim \begin{pmatrix} \parallel & 0 & 0 & \parallel \\ 0 & \perp & \perp & 0 \\ 0 & \perp & \perp & 0 \\ \parallel & 0 & 0 & \parallel \end{pmatrix} \\ &\sim \begin{pmatrix} |\tilde{F}_4|^2 & 0 & 0 & \tilde{F}_4 F_1^*, \tilde{F}_4 F_2^* \\ 0 & |F_1|^2, |F_3|^2 & \text{Re}(F_1 F_3^*) & 0 \\ 0 & \text{Re}(F_1 F_3^*) & |F_1|^2, |F_3|^2 & 0 \\ F_1 \tilde{F}_4^*, F_2 \tilde{F}_4^* & 0 & 0 & |F_1|^2, |F_2|^2, \text{Re}(F_1 F_2^*) \end{pmatrix}. \end{aligned} \quad (25)$$

In the first matrix expression the non-zero elements are indicated with the symbols “ $\parallel$ ” and “ $\perp$ ”. These denote the longitudinal and transverse polarization states, respectively. The second matrix expression indicates the functional dependence on the form factors $F_1(Q^2), \dots, \tilde{F}_4(Q^2)$. Thus, for example, $\widetilde{H}^{03}$ is only non-zero for longitudinally-polarized V 's, and it depends on the structure-function combinations $\tilde{F}_4 F_1^*$ and $\tilde{F}_4 F_2^*$.

Recall that the NP contributions are contained in $\tilde{F}_4(Q^2)$, and that all CP-violating effects require the interference of SM and NP amplitudes. We therefore see immediately that any CP asymmetries will involve $\widetilde{H}^{00}$, $\widetilde{H}^{03}$ or $\widetilde{H}^{30}$. This is consistent with Eqs. (20) and (21), which note that only $\tilde{J}^0$ contains a NP contribution.

By defining symmetric and antisymmetric combinations of the various elements of the tensors $L_{\mu\nu}$ and $\widetilde{H}^{\mu\nu}$, it is possible to write the expression for $L_{\mu\nu} \widetilde{H}^{\mu\nu}$ in a relatively simple manner. Using the notation in Ref. [9],⁶ we write

$$L_{\mu\nu} \widetilde{H}^{\mu\nu} = \sum_X L_X W_X, \quad (26)$$

where $X = A, B, \dots, I, SA, SB, \dots, SG$ and where there is an implied summation over polarization states. Definitions and some expressions for the relevant non-zero contributions are included in the Appendix.

We now examine three possible CP-violating signals: the rate asymmetry, a polarization-dependent rate asymmetry (the polarization is that of the τ), and a triple-product asymmetry.

3.2 Rate Asymmetry

Integrating the differential width in Eq. (23) over the angular variables β and θ , and normalizing the result to the width for $\tau \rightarrow e\nu\bar{\nu}$, we obtain

$$\frac{d\Gamma}{\Gamma_e dQ^2} \simeq \frac{\cos^2 \theta_c}{2 (Q^2)^{3/2} m_\tau^8} (m_\tau^2 - Q^2)^2 |q_1^z| \left[(2Q^2 + m_\tau^2) (W_A + W_B) + 3m_\tau^2 W_{SA} \right], \quad (27)$$

⁶Note the corrections to some of the relevant expressions in the Erratum to Ref. [9].

where $|q_1^z|$ is the magnitude of the momentum of the V in the hadronic rest frame (see the Appendix for an explicit expression) and $\Gamma_e \simeq G_F^2 m_\tau^5 / (192\pi^3)$. Note that W_{SA} contains a Higgs-exchange contribution, but W_A and W_B do not (see the Appendix). In particular,

$$W_{SA} = \widetilde{H}^{00} = \frac{Q^4 (q_1^z)^2}{m_V^2} |\widetilde{F}_4|^2 . \quad (28)$$

The rate asymmetry depends on the difference between the above expression and the analogous one for the τ^+ decay,

$$A_{CP} = \frac{\Delta\Gamma}{\Gamma_{\text{sum}}} , \quad (29)$$

where $\Delta\Gamma$ is the difference of the widths for the process and the anti-process (normalized to Γ_e),

$$\begin{aligned} \Delta\Gamma &= \int \left[\frac{d\Gamma}{\Gamma_e dQ^2} - \frac{d\bar{\Gamma}}{\Gamma_e dQ^2} \right] dQ^2 \\ &\simeq \int \frac{3 \cos^2 \theta_c}{2 (Q^2)^{3/2} m_\tau^6} (m_\tau^2 - Q^2)^2 |q_1^z| (W_{SA} - \overline{W}_{SA}) dQ^2 , \end{aligned} \quad (30)$$

and Γ_{sum} is the sum of the widths for the process and the anti-process.

The expression for $\Delta\Gamma$ may be written explicitly in terms of strong and weak phases. Defining phases in the following manner,

$$F_j(Q^2) = |F_j(Q^2)| \exp [i\delta_j(Q^2)] , \quad (j = 1, \dots, 4) , \quad (31)$$

$$f_H(Q^2) = |f_H(Q^2)| \exp [i\delta_H(Q^2)] , \quad (32)$$

$$b = |b| \exp (i\phi_b) , \quad (33)$$

$$a = |a| \exp (i\phi_a) , \quad (34)$$

we have

$$W_{SA} - \overline{W}_{SA} = 4 \frac{Q^4 (q_1^z)^2}{m_V^2} \frac{|F_4 f_H b|}{m_\tau} \sin (\delta_4 - \delta_H) \sin \phi_b \quad (35)$$

for a vector meson (e.g. $V = \omega$). For an axial-vector meson (e.g. $V = a_1$) the change $b \rightarrow a$ must be made. This expression may be inserted in Eq. (30) to determine $\Delta\Gamma$. The key point is that $\Delta\Gamma$ depends on the product of the NP amplitude (proportional to $f_H b$ or $f_H a$) with the SM scalar form factor F_4 . But F_4 is expected to be very small, regardless of whether V is a vector or axial-vector meson. We therefore conclude that the total rate asymmetries in τ decays are small, even in the presence of NP.

3.3 Polarization-dependent Rate Asymmetry

The regular rate asymmetry defined in Eq. (29) depends on F_4 , and is likely to be very small. It is also possible to define a rate asymmetry that depends on $f_H F_1^*$ and/or $f_H F_2^*$. It turns out that to extract such cross-terms, one need only weight the differential width in Eq. (23) by $\cos \beta$ when performing the integration over $\cos \beta$.⁷ Performing the various integrals and forming a CP asymmetry, we obtain

$$A_{CP}^{\langle \cos \beta \rangle} = \frac{\Delta \Gamma_{\langle \cos \beta \rangle}}{\Gamma_{\text{sum}}} , \quad (36)$$

where $\Delta \Gamma_{\langle \cos \beta \rangle}$ is the difference of the widths for the process and the anti-process:

$$\begin{aligned} \Delta \Gamma_{\langle \cos \beta \rangle} &= \int \left[\frac{d\Gamma}{\Gamma_e dQ^2 d\cos \beta} - \frac{d\bar{\Gamma}}{\Gamma_e dQ^2 d\cos \beta} \right] dQ^2 \cos \beta d\cos \beta \\ &\simeq - \int \frac{\cos^2 \theta_c}{2 (Q^2)^{3/2} m_\tau^6} (m_\tau^2 - Q^2)^2 |q_1^z| (W_{SF} - \bar{W}_{SF}) \rho(Q^2) dQ^2 , \end{aligned} \quad (37)$$

with

$$\rho(Q^2) \equiv \int \left(\cos \psi + P \cos \theta \cos \psi + P \frac{\sqrt{Q^2}}{m_\tau} \sin \theta \sin \psi \right) \frac{d\cos \theta}{2} . \quad (38)$$

Note that ψ is a function of θ (see the Appendix).

We have

$$\begin{aligned} W_{SF} &= \tilde{H}^{03} + \tilde{H}^{30} \\ &= \frac{(Q^2)^{3/2} |q_1^z|}{m_V^2} \left[2 \text{Re} (F_1 \tilde{F}_4^*) \left(\sqrt{Q^2} E_1 \right) + 4 \text{Re} (F_2 \tilde{F}_4^*) |q_1^z|^2 \right] , \end{aligned} \quad (39)$$

where E_1 denotes the energy of the V in the hadronic rest frame. The form factors F_1 and F_2 are expected to be very small for a vector meson (e.g. $V = \omega$) but are large for an axial-vector meson (e.g. $V = a_1$). We therefore conclude that this asymmetry can be sizeable for the decay $\tau \rightarrow a_1 \pi \nu_\tau$.

Figure 2 shows plots of $\rho(Q^2)$ for a few parameter choices, assuming that the τ 's are produced in a symmetric collider environment. We see that $\rho(Q^2)$ can be large, even if the τ 's are unpolarized ($P = 0$).

Employing the definitions in Eqs. (31)-(34), and assuming that the decay is $\tau \rightarrow a_1 \pi \nu_\tau$, we find the following expression for the difference $W_{SF} - \bar{W}_{SF}$,

$$\begin{aligned} W_{SF} - \bar{W}_{SF} &= 4 \frac{(Q^2)^{3/2} |q_1^z|}{m_{a_1}^2} \frac{|f_H a|}{m_\tau} \\ &\times \left[|F_1| \sin(\delta_1 - \delta_H) \sqrt{Q^2} E_1 + 2 |F_2| \sin(\delta_2 - \delta_H) (q_1^z)^2 \right] \sin \phi_a, \end{aligned} \quad (40)$$

⁷Ref. [7] shows some partial results along these lines, but assumes that the scalar contributions are zero. Including the scalar contributions is straightforward.




Figure 2: Plots of $\rho(Q^2)$ for various values of the τ polarization P . The function $\rho(Q^2)$ is involved in the polarization-dependent CP asymmetry and is defined in Eq. (38). The solid and dashed lines show the results for $\sqrt{s} = m_{\Upsilon(4S)}$ and $\sqrt{s} = m_Z$, respectively.

This expression may be inserted in Eq. (37) to determine $\Delta\Gamma_{\langle\cos\beta\rangle}$. We stress that the angle β must be measured in this CP asymmetry. That is, in contrast to the total rate asymmetry (which is very small), the polarization-dependent rate asymmetry is in fact a *directional* asymmetry.

As noted above, the four π 's in $\tau \rightarrow \pi\pi\pi\pi\nu_\tau$ come primarily from the decay of $\omega\pi$ or $a_1\pi$. The polarization-dependent rate asymmetry of $\omega\pi$ is essentially zero, so the (inclusive) measurement of this asymmetry in $\tau \rightarrow \pi\pi\pi\pi\nu_\tau$ effectively probes the (exclusive) contribution of $\tau \rightarrow a_1\pi\nu_\tau$ and it appears that one does not require that the a_1 be reconstructed. However, there is a potential problem here. The angle β is the angle between the a_1 momentum in the hadronic rest frame and the direction of the lab as viewed in this frame. In order to get β , one does in fact have to reconstruct the a_1 to get its momentum. However, the width of the a_1 is extremely large. As a consequence, it may not be possible to identify unambiguously which three of the four pions came from the a_1 . If so, there will be an ambiguity in defining β . There are two possible solutions to this problem. One is to exclude events in which one cannot unambiguously reconstruct the a_1 . This may severely reduce the number of events. The second is to include all possible reconstructions of the a_1 [15]. This may severely reduce the signal. We assume that the experimentalists will find the optimal solution.

We now present a brief numerical analysis of the polarization-dependent asymmetry for $\tau \rightarrow a_1\pi\nu_\tau$. The polarization asymmetry depends on cross terms involving the NP form factor f_H with the SM form factors F_1 and F_2 [see Eq. (40)]. The form

factors F_1 and F_2 are expected to dominate in the calculation of the width for $\tau \rightarrow a_1 \pi \nu_\tau$, but at this point the relative sizes of the two are not known. To perform a few estimates of the polarization asymmetry, we make the simplifying assumption that either $F_1 = 0$ or $F_2 = 0$.⁸ In addition, since the Q^2 dependence of the form factors is theoretically unknown, we have repeated the calculations for two different sets of assumptions in each case.

In the first scenario, we assume that F_1 (or F_2) receives contributions only from the $\rho(770)$; in the second, we assume that both the $\rho(770)$ and $\rho(1450)$ resonances contribute. The form factors thus have the form shown for F_3^{pole} in Eq. (14), but with $g_{\rho\omega\pi}/\gamma_\rho$ replaced by a constant that is determined by the experimental rate for $\mathcal{B}(\tau \rightarrow a_1 \pi \nu_\tau)$.⁹ The CLEO collaboration has performed several fits to determine the branching ratio for $\tau \rightarrow a_1 \pi \nu_\tau$. In one of the better fits, the collaboration obtains $R_{a_1\pi} = \mathcal{B}(\tau^- \rightarrow (a_1\pi)^- \nu_\tau) / \mathcal{B}(\tau^- \rightarrow 2\pi^- \pi^+ \pi^0 \nu_\tau) = 0.49 \pm 0.02 \pm 0.02$ (Model 3 in Table IV of Ref. [15]). There is a slight subtlety in that this ratio includes contributions from both $\tau^- \rightarrow a_1^0 \pi^- \nu_\tau$ and $\tau^- \rightarrow a_1^- \pi^0 \nu_\tau$. Furthermore, the a_1^0 can be formed in two different ways, while the a_1^- can only be formed in one way. The experimental analysis includes all three possibilities and even allows them, in principle, to interfere. For our analysis we consider the a_1^0 case and, as an approximation, use the central value $R_{a_1^0\pi^-} \simeq (2/3) \times R_{a_1\pi} \simeq 0.327$ to determine F_1 (or F_2). We then assume that a NP contribution could reasonably be assumed to be “hidden” in the remaining uncertainty. We thus tune the NP combination $|af_H|$ such that the NP contribution to $R_{a_1^0\pi^-}$ is equal to approximately 0.019 (i.e., we multiply the uncertainty by 2/3 as well). This procedure yields the value $|af_H| \simeq 6.8$. (Note that f_H is assumed to be a constant with zero strong phase.) Having fixed F_1 (or F_2) in this manner, and setting $P=1$ in Eq. (38), we proceed to compute the polarization asymmetry defined in Eq. (36).

If $F_{1,2}$ are assumed to be dominated by the $\rho(770)$ resonance we obtain quite small asymmetries – of order $\sim 1\%$ for $|\sin \phi_a| = 1$ – regardless of which form factor has been set to zero. This fact is not surprising in this case: since the $\rho(770)$ cannot go on shell in $\tau \rightarrow a_1 \pi \nu_\tau$, the factors $\sin(\delta_{1,2} - \delta_H)$ in Eq. (40) are small (recall that we have assumed $\delta_H = 0$). Larger asymmetries are obtained by assuming that both the ρ and ρ' contribute to $F_{1,2}$, since the ρ' can go on-shell, leading to appreciable strong phase effects. In this case asymmetries of order 15% (7.5%) can be obtained when F_1 (F_2) is set to zero (again assuming that the strong phase associated with the NP contribution, δ_H , is set to zero).

⁸Although the real picture is expected to correspond to some intermediate situation, this seems to be a reasonable way to proceed in order to make model-independent predictions.

⁹In the first case we set $A_1 = 0$ (i.e., ρ resonance only). In the second case we somewhat arbitrarily take $A_1 = -0.24$, which is the value we also use in Sec. 3.4 when estimating the triple product asymmetry in $\tau \rightarrow \omega \pi \nu_\tau$. In that case the choice is well-motivated [15].

3.4 Triple-product Asymmetry

If the momentum of the τ can be determined experimentally, it is possible to construct a triple-product CP asymmetry. In this subsection we will assume that the τ is unpolarized ($P = 0$), but that the polarization of the V is measured. Furthermore, we will assume that the polarization of the V is neither purely transverse, nor purely longitudinal. As we shall see, this assumption will allow the NP contribution to interfere with the SM form factor F_3 , allowing for possibly observable effects for $V = \omega$ and $V = \rho$.

There are in principle two possible sources of triple products in hadronic τ decays, assuming that the τ 's are unpolarized. The first possible source arises from the leptonic tensor $L_{\mu\nu}$, which contains a term proportional to $i\varepsilon_{\alpha\mu\beta\nu}l'^\alpha l^\beta$. This term is contracted with $\tilde{H}^{\mu\nu} = \tilde{J}^\mu (\tilde{J}^\nu)^\dagger$, which only contains NP contributions if μ and/or ν is equal to zero [see Eqs. (20) and (21)]. It is then evident that the $i\varepsilon_{\alpha\mu\beta\nu}l'^\alpha l^\beta$ term in $L_{\mu\nu}$ cannot lead to a CP-odd TP asymmetry. (Note that $i\varepsilon_{i0jk}l'^i l^j = 0$ because $\vec{l}' = \vec{l}$ in the hadronic rest frame.)

The only other source of a CP-odd TP asymmetry is found in the hadronic tensor; the triple product arises from a Levi-Civita tensor term in $\tilde{J}^\mu$ involving the F_3 term. This F_3 can be large if V is a vector meson: $V = \omega, \rho$. The CP-violating TP asymmetry can be sizeable for the decays $\tau \rightarrow \omega\pi\nu_\tau$ and $\tau \rightarrow \rho\pi\nu_\tau$.

The TP involves the V momentum, so that it is necessary to reconstruct the V . This makes the decay $\tau \rightarrow \rho\pi\nu_\tau$ problematic. The width of the ρ is very large and one runs into a similar problem as that described in Sec. 3.3. In the decay $\tau \rightarrow \rho\pi\nu_\tau \rightarrow 3\pi\nu_\tau$, it is not possible to identify unambiguously which two of the three pions came from the ρ . It is therefore very difficult to reconstruct the ρ and, as a consequence, difficult to measure the TP in these decays.

The decay $\tau \rightarrow \omega\pi\nu_\tau$ does not suffer the same problem. The ω is quite narrow, so it can be fairly easily reconstructed. For this reason, in the measurement of the TP asymmetry, we exclude $\tau \rightarrow \rho\pi\nu_\tau$, concentrating only on $\tau \rightarrow \omega\pi\nu_\tau$. (Note that this implies that there are no clean CP-violating signals in $\tau \rightarrow 3\pi\nu_\tau$.)

Keeping only the terms that can lead to a CP-odd TP asymmetry, and averaging over the spin states of the initial τ , we have

$$L_{\mu\nu}\tilde{H}^{\mu\nu}|_{\text{TP}} = \left[-L^{0i}\tilde{J}^0 (J^i)^\dagger - L^{i0}J^i (\tilde{J}^0)^\dagger \right]_{\text{TP}} \quad (41)$$

$$= \frac{8m_\tau Q^2}{E_1} \text{Im}(bf_H F_3^*) (\vec{\epsilon}_1 \cdot \vec{q}_1) \vec{\epsilon}_1 \cdot (\vec{l} \times \vec{q}_1) . \quad (42)$$

In the second line we have adopted a real basis for the ω polarization tensor and have used the relation $\epsilon_1 \cdot q_2 = \sqrt{Q^2} \vec{\epsilon}_1 \cdot \vec{q}_1 / E_1$, which is valid in the hadronic rest frame.

Since we now assume that the τ direction is known, we can simplify our coordinate system somewhat compared to that shown in Fig. 1. We define ζ to be the

angle between $\vec{l}$ and $\vec{q}_1$ ($\vec{l}$ is the momentum of the τ ; $\vec{q}_1$ is the momentum of the V); that is,

$$\vec{l} \cdot \vec{q}_1 = |\vec{l}| |\vec{q}_1| \cos \zeta . \quad (43)$$

It is then convenient to define three direction vectors as follows,

$$\vec{n}_1 = \frac{\vec{q}_1}{|\vec{q}_1|} \frac{m_\omega}{E_1} , \quad \vec{n}_2 = \frac{\vec{l} \times \vec{q}_1}{|\vec{l} \times \vec{q}_1|} = \frac{\vec{l} \times \vec{q}_1}{|\vec{l}| |\vec{q}_1| \sin \zeta} , \quad \vec{n}_3 = \frac{(\vec{l} \times \vec{q}_1) \times \vec{q}_1}{|(\vec{l} \times \vec{q}_1) \times \vec{q}_1|} , \quad (44)$$

where all quantities are expressed in the hadronic rest frame. The peculiar normalization of $\vec{n}_1$ is associated with the boost from the ω 's rest frame to the hadronic rest frame. With the above definitions, the condition $\epsilon_1^2 = -1$ becomes

$$(\vec{\epsilon}_1 \cdot \vec{n}_1)^2 + (\vec{\epsilon}_1 \cdot \vec{n}_2)^2 + (\vec{\epsilon}_1 \cdot \vec{n}_3)^2 = 1 . \quad (45)$$

Rewriting Eq. (42) in terms of the various direction vectors, we have

$$L_{\mu\nu} \widetilde{H}^{\mu\nu} \Big|_{\text{TP}} = \frac{8m_\tau |q_1^z| Q^2}{m_\omega} \sin \zeta \text{Im} (bf_H F_3^*) (\vec{\epsilon}_1 \cdot \vec{n}_1) (\vec{\epsilon}_1 \cdot \vec{n}_2) . \quad (46)$$

Substituting this expression into Eq. (17) and calculating the differential width, we find,

$$\frac{d\Gamma}{\Gamma_e dQ^2} \Big|_{\text{TP}} = \frac{3\pi \cos^2 \theta_c}{4 m_\omega m_\tau^6 \sqrt{Q^2}} (m_\tau^2 - Q^2)^2 |q_1^z|^3 \text{Im} (bf_H F_3^*) (\vec{\epsilon}_1 \cdot \vec{n}_1) (\vec{\epsilon}_1 \cdot \vec{n}_2) , \quad (47)$$

in which we have integrated over the angle ζ .

It is now possible to define a CP asymmetry based on Eq. (47),

$$A_{CP}^{\text{TP}} = \frac{\Delta\Gamma_{\text{TP}}}{\Gamma_{\text{sum}}} , \quad (48)$$

where $\Delta\Gamma_{\text{TP}}$ is the difference of the widths for the process and the anti-process,

$$\begin{aligned} \Delta\Gamma_{\text{TP}} &= \int \left[\frac{d\Gamma}{\Gamma_e dQ^2} - \frac{d\bar{\Gamma}}{\Gamma_e dQ^2} \right]_{\text{TP}} dQ^2 \\ &\simeq (\vec{\epsilon}_1 \cdot \vec{n}_1) (\vec{\epsilon}_1 \cdot \vec{n}_2) \int \frac{3\pi \cos^2 \theta_c}{2 m_\omega m_\tau^6 \sqrt{Q^2}} (m_\tau^2 - Q^2)^2 |q_1^z|^3 \\ &\quad \times |bf_H F_3| \cos(\delta_3 - \delta_H) \sin \phi_b dQ^2 . \end{aligned} \quad (49)$$

This CP-odd TP asymmetry depends on the form-factor combination $f_H F_3$, which is the most favourable combination possible, since F_3 is known to dominate over the other form factors in Eq. (11) (for the decay $\tau \rightarrow \omega \pi \nu_\tau$). Furthermore, as noted

above, the asymmetry vanishes if the ω 's polarization is either purely longitudinal ($\vec{\epsilon}_1 \cdot \vec{n}_2 = 0$) or purely transverse ($\vec{\epsilon}_1 \cdot \vec{n}_1 = 0$). Finally, note that the TP asymmetry depends on the cosine of the strong phase difference, so it can be non-zero even if there are no strong phases.

The TP asymmetry defined above is written explicitly in terms of projections of the polarization vector onto the vectors $\vec{n}_1$ and $\vec{n}_2$. A similar approach was followed in Refs. [18, 19]. In that case the authors constructed T-odd observables in $B \rightarrow D^* \ell \nu$ using the polarization of the D^* . In Appendix B of Ref. [19], the authors showed that there was a direct connection between the asymmetries written in terms of polarization projections and more “physical” asymmetries constructed using the momenta of the D^* decay products.

We conclude this section by presenting a numerical investigation of the size of the TP asymmetry in the decay $\tau \rightarrow \omega \pi \nu_\tau$. This asymmetry is easier to study than the one discussed in the previous section, since the SM decay amplitude is dominated by a single SM form factor, F_3 . For our numerical estimate of the TP asymmetry, we use the expression for F_3 given in Eq. (14), taking $A_1 = -0.24$, $g_{\rho\omega\pi} = 16.1 \text{ GeV}^{-1}$ and $\gamma_\rho \simeq 4.95$ (Model 2 in Table I of Ref. [15]). This yields $R_{\omega\pi} = \mathcal{B}(\tau^- \rightarrow (\omega\pi)^- \nu_\tau) / \mathcal{B}(\tau^- \rightarrow 2\pi^- \pi^+ \pi^0 \nu_\tau) \simeq 0.38$, in agreement with Model 3 of Table IV of Ref. [15], which had obtained the value $R_{\omega\pi} = 0.38 \pm 0.02 \pm 0.01$. Assuming that the NP contribution can at most be responsible for the remaining uncertainty in $R_{\omega\pi}$, we tune $|bf_H|$ such that its contribution to $R_{\omega\pi}$ is approximately 0.022. This procedure yields $|bf_H| \simeq 0.98$. Finally, we have computed the asymmetry using this value and assuming that the strong phase $\delta_H = 0$ and that $|\sin \phi_b| = 1$. The resulting TP asymmetry has a magnitude of order 30% multiplied by $(\vec{\epsilon}_1 \cdot \vec{n}_1)(\vec{\epsilon}_1 \cdot \vec{n}_2)$

4 Conclusions

In this paper we have re-examined CP violation in $\Delta S = 0$ hadronic τ decays. Any CP-violating effect requires the interference of (at least) two amplitudes with a relative weak phase (and often a relative strong phase). The standard-model contribution involves (essentially) only a virtual W and constitutes one amplitude. The second amplitude must come from new physics (NP). We assume that the NP consists of a charged Higgs boson. This is a fairly general assumption.

However, in many models, the coupling of the charged Higgs boson to first-generation final states is proportional to first-generation masses, and is negligible. In order to obtain a measurable CP-violating effect in τ decays, we must examine non-“standard” NP, and consider only the case in which the charged-Higgs coupling to first-generation particles is large. This is not usual, but if CP violation is seen in τ decays, it points clearly to a different type of NP.

The decays examined are essentially $\tau \rightarrow N\pi\nu_\tau$, with $N = 2, 3, 4$. We have shown that the charged Higgs H does not couple to two pions, and so the decay $\tau \rightarrow \pi\pi\nu_\tau$ will not exhibit CP violation. However, H does couple to 3π and 4π , so that CP

violation could be seen in $\tau \rightarrow N\pi\nu_\tau$ ($N = 3, 4$). In addition to non-resonant contributions, these decays receive resonant contributions from intermediate particles such as ω , ρ and a_1 . These resonant contributions involve a smaller number of final particles, and are dominant. We therefore consider only the resonant contributions to $\Delta S = 0$ hadronic τ decays.

For the decay $\tau \rightarrow V\pi\nu_\tau$ (V is a vector or axial-vector meson), the SM hadronic current involves four form factors $F_{1-4}(Q^2)$. If V is a vector meson (ω , ρ), the contribution proportional to F_3 dominates. The F_1 and F_2 terms violate G -parity, and there is no evidence for such second-class currents. Equally, there is no evidence for a non-zero F_4 term, which is scalar. If V is an axial-vector meson (a_1), F_1/F_2 and F_3 change roles. Thus, in the decay $\tau \rightarrow a_1\pi\nu_\tau$, the contributions proportional to F_1 and F_2 dominate, while the F_3 term violates G -parity. For vector mesons, the NP hadronic current is proportional to bf_H , where f_H is a form factor and b is assumed to have a non-zero weak phase (b is the pseudoscalar coupling of the charged Higgs). For axial-vector mesons, one makes the substitution $b \rightarrow a$, where a is the scalar coupling of the charged Higgs.

We have examined three potential signals of CP violation: an overall rate asymmetry, a polarization-dependent rate asymmetry (the polarization is that of the τ), and a triple-product asymmetry. It is found that the overall rate asymmetry is proportional to $|f_H F_4|$ and is tiny for all hadronic τ decays.

The polarization-dependent rate asymmetry is proportional to $|f_H F_1|$ and $|f_H F_2|$. It is therefore sizeable only for $\tau \rightarrow a_1\pi\nu_\tau$. However, the width of the a_1 is extremely large, which can lead to ambiguities in reconstructing the direction of the a_1 momentum. If this problem can be addressed, the polarization-dependent rate asymmetry in $\tau \rightarrow a_1\pi\nu_\tau \rightarrow 4\pi\nu_\tau$ is a good signal of CP violation, with values of order 15% currently allowed by experimental data.

Finally, the triple-product asymmetry is proportional to $|f_H F_3|$, and can be sizeable for $\tau \rightarrow \omega\pi\nu_\tau$ and $\tau \rightarrow \rho\pi\nu_\tau$. However, the ρ width is very large, which can lead to difficulties in reconstructing the ρ from $\tau \rightarrow 3\pi\nu_\tau$. The ω is quite narrow and does not suffer similar problems. For this reason we consider only the decay $\tau \rightarrow \omega\pi\nu_\tau$ in the calculation of the TP asymmetry. Note that this means that there are no clean signals of CP violation in $\tau \rightarrow 3\pi\nu_\tau$. Our analysis indicates that the current experimental data allow a TP asymmetry for $\tau \rightarrow \omega\pi\nu_\tau$ as large as 30% multiplied by $(\vec{\epsilon}_1 \cdot \vec{n}_1)(\vec{\epsilon}_1 \cdot \vec{n}_2)$.

Note also that this triple product is $\epsilon_\omega \cdot (p_\omega \times p_\tau)$. In order to measure this TP, one needs the spin of the ω (ϵ_ω), which in turn means that the direction of the τ must be known.

Acknowledgments: We thank the following people for helpful communications: S. Banerjee, T. Browder, H. Hayashii, B. Kowalewski, M. Roney and B. Viaud. A.D. and K.K. thank the Physics Departments of the University of Toronto and the Université de Montréal, respectively, for their hospitality during the initial stages of this work. This work was financially supported by NSERC of Canada. The work of

K.K. was supported in part by the U.S. National Science Foundation under Grants PHY-0301964 and PHY-0601103.

A Appendix

This section contains the definitions for the relevant L_X and W_X in Eq. (26) as well as explicit expressions for several other quantities, such as $|q_1^z|$, $\cos \theta$ and $\cos \psi$.

A.1 Expressions for L_X and W_X .

In the polarization basis indicated in Eqs. (24) many of the elements of the hadronic tensor $\widetilde{H}^{\mu\nu}$ are zero. Summing over the polarization states, we find

$$L_{\mu\nu}\widetilde{H}^{\mu\nu} = L_A W_A + L_B W_B + L_E W_E + L_{SA} W_{SA} + L_{SF} W_{SF} + L_{SG} W_{SG} , \quad (50)$$

where [9],

$$\begin{aligned} L_A &= \frac{1}{2} (L^{11} + L^{22}) , & W_A &= \widetilde{H}^{11} + \widetilde{H}^{22} , \\ L_B &= L^{33} , & W_B &= \widetilde{H}^{33} , \\ L_E &= \frac{i}{2} (L^{12} - L^{21}) , & W_E &= -i (\widetilde{H}^{12} - \widetilde{H}^{21}) , \\ L_{SA} &= L^{00} , & W_{SA} &= \widetilde{H}^{00} , \\ L_{SF} &= -\frac{1}{2} (L^{03} + L^{30}) , & W_{SF} &= \widetilde{H}^{03} + \widetilde{H}^{30} , \\ L_{SG} &= -\frac{i}{2} (L^{03} - L^{30}) , & W_{SG} &= -i (\widetilde{H}^{03} - \widetilde{H}^{30}) . \end{aligned}$$

Explicit expressions for these quantities may be found in Ref. [7] (recalling that one needs to make the replacement $F_4 \rightarrow \widetilde{F}_4$). A few expressions that are useful for our purposes are the following,

$$W_A = 2Q^2 [(q_1^z)^2 |F_3|^2 + Q^2 |F_1|^2] , \quad (51)$$

$$W_B = \frac{Q^2}{m_V^2} \left[Q^2 E_1^2 |F_1|^2 + 4\sqrt{Q^2} E_1 (q_1^z)^2 \text{Re}(F_1 F_2^*) + 4 (q_1^z)^4 |F_2|^2 \right] , \quad (52)$$

$$W_{SA} = \frac{Q^4 (q_1^z)^2}{m_V^2} |\widetilde{F}_4|^2 , \quad (53)$$

$$W_{SF} = \frac{(Q^2)^{3/2} |q_1^z|}{m_V^2} \left[2 \text{Re}(F_1 \widetilde{F}_4^*) \left(\sqrt{Q^2} E_1 \right) + 4 \text{Re}(F_2 \widetilde{F}_4^*) |q_1^z|^2 \right] , \quad (54)$$

$$W_{SG} = \frac{(Q^2)^{3/2} |q_1^z|}{m_V^2} \left[2 \text{Im}(F_1 \widetilde{F}_4^*) \left(\sqrt{Q^2} E_1 \right) + 4 \text{Im}(F_2 \widetilde{F}_4^*) |q_1^z|^2 \right] . \quad (55)$$

Note that only W_A is non-zero in the limit that the F_3 structure function dominates the current $\tilde{J}^\mu$, as is the case for $\tau \rightarrow \omega\pi\nu_\tau$. Furthermore, as noted in the text, the form factor $\tilde{F}_4$, which contains the Higgs contribution, only appears in combination with F_1 , F_2 and itself, and not with the dominant F_3 term. The combination $F_3\tilde{F}_4^*$ can arise if the V 's polarization is taken to have non-zero projections in both the transverse and longitudinal directions. Such is the case for the triple-product CP asymmetry constructed in the text.

A.2 Expressions for some kinematical quantities.

In the hadronic rest frame the magnitude of the momentum of the V is given by

$$|q_1^z| = \frac{1}{2\sqrt{Q^2}} \lambda^{1/2}(Q^2, m_V^2, m_\pi^2) , \quad (56)$$

where $\lambda(x, y, z) \equiv x^2 + y^2 + z^2 - 2(xy + xz + yz)$. Then the energy of the V is simply $E_1 = \sqrt{(q_1^z)^2 + m_V^2}$.

It is also possible to derive a relation between the angles θ and ψ . Assuming that the τ 's are pair-produced at a symmetric collider, these angles may be expressed as follows [7],

$$\cos \theta = \frac{2xm_\tau^2 - m_\tau^2 - Q^2}{(m_\tau^2 - Q^2)(1 - 4m_\tau^2/s)^{1/2}} , \quad (57)$$

$$\cos \psi = \frac{x(m_\tau^2 + Q^2) - 2Q^2}{(m_\tau^2 - Q^2)(x^2 - 4Q^2/s)^{1/2}} , \quad (58)$$

where $s = 4E_{\text{beam}}^2 = 4E_\tau^2$ and $x = 2E_h/\sqrt{s}$, with E_h being the hadron energy in the lab frame. To determine the relation between ψ and θ explicitly, one can solve Eq. (57) for x in terms of $\cos \theta$ and then substitute the result into Eq. (58). This allows one, for example, to perform the integration in Eq. (38). Similar expressions could be derived in the case of an asymmetric collider, assuming one knew the correct boost factor.

In determining the polarization-dependent CP asymmetry, we have assumed the incident τ 's to have polarization P as viewed from the lab. A derivation of the τ spin four-vector in the hadronic rest frame may be found in Appendix A of Ref. [9]. For convenience, we list the result here as well. In the S' frame (see Fig. 1),

$$(s^\mu)^{S'} = P \left(-\frac{|\vec{l}|}{m_\tau} \cos \theta, -\frac{l_0}{m_\tau} \cos \theta \sin \psi + \sin \theta \cos \psi, 0, \right. \\ \left. -\frac{l_0}{m_\tau} \cos \theta \cos \psi - \sin \theta \sin \psi \right) , \quad (59)$$

where l_0 and $|\vec{l}|$ refer to the energy and momentum of the τ in the hadronic rest frame. To obtain the expression in the S frame, one needs to perform an orthogonal rotation on the spatial components of $(s^\mu)^{S'}$ using the angles α and β indicated in Fig. 1.

References

- [1] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592**, 1 (2004).
- [2] S. Y. Choi, J. Lee and J. Song, Phys. Lett. B **437**, 191 (1998);
- [3] J. H. Kuhn and E. Mirkes, Phys. Lett. B **398**, 407 (1997).
- [4] S. Y. Choi, K. Hagiwara and M. Tanabashi, Phys. Rev. D **52**, 1614 (1995).
- [5] Y. S. Tsai, arXiv:hep-ph/9801274.
- [6] R. Decker, Z. Phys. C **36**, 487 (1987).
- [7] R. Decker and E. Mirkes, Z. Phys. C **57**, 495 (1993).
- [8] H. Davoudiasl and M. B. Wise, Phys. Rev. D **53**, 2523 (1996).
- [9] J. H. Kuhn and E. Mirkes, Z. Phys. C **56**, 661 (1992) [Erratum-ibid. C **67**, 364 (1995)].
- [10] J. H. Kuhn and E. Mirkes, Phys. Lett. B **398**, 407 (1997).
- [11] J. H. Kuhn, Phys. Lett. B **313**, 458 (1993).
- [12] P. Avery *et al.* [CLEO Collaboration], Phys. Rev. D **64**, 092005 (2001).
- [13] I. I. Bigi and A. I. Sanda, Phys. Lett. B **625**, 47 (2005).
- [14] G. Bonvicini *et al.* [CLEO Collaboration], Phys. Rev. Lett. **88**, 111803 (2002);
S. Anderson *et al.* [CLEO Collaboration], Phys. Rev. Lett. **81**, 3823 (1998).
- [15] K. W. Edwards *et al.* [CLEO Collaboration], Phys. Rev. D **61**, 072003 (2000).
- [16] P. Bourdon, Nucl. Phys. Proc. Suppl. **40**, 203 (1995).
- [17] D. Buskulic *et al.* [ALEPH Collaboration], Z. Phys. C **74**, 263 (1997).
- [18] G. H. Wu, K. Kiers and J. N. Ng, Phys. Lett. B **402**, 159 (1997).
- [19] G. H. Wu, K. Kiers and J. N. Ng, Phys. Rev. D **56**, 5413 (1997).